

XXV International Population Conference
Tours, France
Paper for Session 112
“Biomarkers, health and demographic research”, July 21, 2005

A draft outline. Please, do not cite or quote.

OBJECTIVE SLEEP DURATION AND HEALTH IN ELDERLY RUSSIANS

Maria Shkolnikova¹, Blake Aber², Maxine Weinstein³, Lyubov' Kravtsova¹, Svetlana Shalnova⁴, Vladimir Shkolnikov⁵, and James W. Vaupel⁵

E-mail address for communication: shkolnikov@demogr.mpg.de

¹ Center for Cardiac Arrhythmia, Moscow Institute of Pediatrics and Child Surgery;

² Columbia University, New York;

³ Georgetown University, Washington DC;

⁴ State Research Centre for Preventive Medicine, Moscow;

⁵ Max Planck Institute for Demographic Research, Rostock, Germany.

ABSTRACT

It is the first examination of relationship between sleep duration and health in the high mortality context of Russia. Night and daytime sleep durations are measured from self-reporting and from 24 hour heart rate trends (Holter monitoring). The sample of 201 individuals is drawn from the Moscow Lipid Research Clinics (LRC) cohort, followed up since 1975-77. Field works were conducted in 2002-03. Holter data are available for 185 individuals. Although objective and reported mean sleep durations are nearly the same (8 h 14 min and 8 h 07 min), there are significant intra-individual differences. OLS and ordered logit regressions controlling for age and sex reveal significant associations between objective sleep (night and night+day sleep durations, and presence of daytime sleep) and health outcomes. No significant relationships with health outcomes are found for subjective night sleep. For objective sleep, longer sleep corresponds to lower grip strength, poorer self-rated health, poorer immediate recall, and a higher mortality risk score (based on the past mortality experience of the LRC cohort). Long sleep duration is a marker of ill-health and, potentially, mortality. Predictive power increases if objective sleep measures are used, a consideration which is especially important in small-size studies.

Keywords: heart rate, monitoring, Holter, mortality, aging

The relationship between length of sleep and health has been studied in some detail. Analyses of three large follow-up studies have provided compelling evidence of higher mortality risk associated with both short and long sleep durations (1, 2, 3). A summary paper that reviews 18 earlier studies on sleep and health suggests a generally U-shaped relationship between sleep duration and mortality, with long sleep being of more concern than short sleep (4). Elevated mortality risk was found even among persons sleeping a “normal” 8-8.5 hours, compared with those sleeping 7-7.5 hours. Long sleep durations have been linked to higher risk for diabetes (5), heart disease (2), cancer (6), and stroke (7). Because large fractions of the population (25 to 50 percent of people in 7 European countries sleep eight hours or more (8)), these results suggest a potentially high population-attributable risk of longer sleep durations.

Sleep deprivation has generally been found to have a negative effect on cognition (9, 10), but little is known about the effects of long sleep duration on cognition. The relation between sleep and muscular function is also unclear. Some studies have found that sleep deprivation affects endurance

(11, 12). However, another study found no association between sleep duration and the strength of various muscle groups (13).

Most studies use self-reported data on when the respondents go to sleep and when they awaken, however, respondents often report inaccurate information. Important differences between self-reported and polysomnography measurements have been identified (22, 23, 24). Several authors have noted the subjective nature of, and significant errors in, self-reported sleep data (25, 26, 4).

This is the first epidemiological study linking sleep duration with health outcomes in elderly Russians. The Russian situation is of particular interest: During the last three decades this country has been facing an increasing burden of disease and mortality (29). Many scholars connect the Russian health crisis with psychosocial factors (30), which can be related to sleep (31).

This study uses six measures of overall health status: number of mobility limitations, grip strength, immediate recall, self-reported health, perceived stress, and a composite mortality risk score based on risk factors measured approximately 25 years earlier.

MATERIALS AND METHODS

We use data collected by the Moscow Pilot Project on “Biological mechanisms of stress-related hazard in Russia,” jointly conducted by the Max Planck Institute of Demographic Research (Rostock, Germany) and the State Research Centre for Preventive Medicine (Moscow). Participants were selected randomly from the survivors of the Moscow Lipid Research Clinic (LRC) cohort. The LRC cohort was originally established in 1975-77 and has served as a central source of epidemiological information about the Russian population for the last two decades (31, 33, 34, 35, 36, 37). 24-hour Holter monitoring was performed by two-channel tape recorders (Schiller AG, Switzerland) according to the guidelines of the American Heart Association (38).

The study subjects were asked to complete diaries which contained the starting and ending times of night and daytime sleep. These data were used for calculation of reported sleep durations. Sleep and activity periods were also identified from the 24-hour heart rate trends based on the original Holter monitor heart rate recordings (Figure 1). HR trends were constructed using one-minute averaging of continuous HR with calculation of maximum, minimum, and mean values. This method allows detection of changes in HR trends related to sleep with high precision.

The onset of night sleep can be seen on a HR trend from a sharp decrease in the heart rate followed by a plateau of a long-standing low heart rate (Figure 1). Awakening is identified as a reverse transition from the plateau to a higher heart rate with higher amplitude of fluctuations. Short-term awake periods can be seen on the HR trends (Figure 1).

In our analyses sleep is described by: subjective night sleep duration (from the diary), objective night sleep duration, presence of daytime sleep, and total (day plus night) sleep duration. The last three variables were based on Holter data. Our analyses treat hours of objective and reported night sleep as continuous variables. Daytime sleep was observed in only 39 subjects and its duration was highly variable. It was included as an additive component into the continuous total sleep duration. It was also treated as a dichotomous variable in some regressions (“presence” of daytime sleep).




Figure 1. Example of diurnal and nocturnal heart rate dynamics (duration = 21 h 40 min). HR daytime average = 82 beat/min, HR night average = 58 beat/min, night sleep duration = 8 h 50 min, ↓ – beginning of night sleep, ↑ – end of night sleep. Moscow Pilot Project “Biological mechanisms of stress-related hazard in Russia”, 2002-03.

We used standard regression techniques to examine the relation between each of the six health outcomes and sleep. Self-rated health was modeled as an ordered logit; the other five health outcomes were modeled using ordinary least squares. All analyses were done using STATA 7.0 (43).

RESULTS

Table 3 shows several characteristics of the study sample and of sleep.

All study subjects were retired and living at home in their family context. Average duration of reported night sleep was 494 min (8 h 13 min) with a range of 3 to 12 hours. According to diaries, on average the subject’s sleep began at 23 h 07 min. The average time of awakening was 7 h 20 min.

On average, objective night sleep duration was 487 min (8 h 07 min) with a range of 4 to 13 hours. Median duration of night sleep was 8 h 10 min. Five percent of the sample slept less than 5 h 50 min and five percent slept more than 10 h 10 min. On average, night sleep began at 23 h 09 min with 5-th and 95-th percentiles being 21 h 20 min and 01 h 20 min, respectively. The average time of awakening was 7 h 16 min with 5-th and 95-th percentiles at 5 h 20 min and 9 h 17 min.

Regressions of sleep duration on sex and age showed no significant difference between men and women and no significant trend across ages. Thus, the average values of the objective and subjective night sleep durations are almost the same. The two measures have a correlation coefficient of 0.69 ($p=0.000$). However, individual differences between the two durations are substantial: 47 percent of reported durations differed from the objective ones by more than 30 min. There was no evidence of a consistent bias toward either over- or under-reporting duration. Reported night sleep durations show considerable heaping on integer values: 51 percent of individual sleep durations are 7, 8, 9, or 10 hours, with 18 percent of this majority reporting a sleep duration of 8 hours. Respondents clearly tended to round times of beginning and end of the night sleep resulting in integer durations. By contrast, round numbers of hours constituted only 13 percent of the objective sleep durations.

Daytime sleep episodes were found in 39 individuals (21 percent); mean duration was 91 min and ranged from 20 to 210 min.

Table 4 shows the results of the regressions of the six health outcomes on sleep characteristics. Every combination of a health outcome with a sleep variable corresponds to one regression and includes controls for age and sex. The rows of Table 4 present five regression models. The first row shows results from regressions of each health outcome (shown in the columns) on reported night sleep duration; it is comparable to many other studies. The second regresses each outcome on objective night sleep duration. The third regresses each outcome on objective night sleep duration, but with a restricted set of observations (six individuals with night sleep durations of 5.5 hours or

less excluded). We exclude these six cases because our data do not provide adequate power for us to model accurately the relationship between short sleep and health outcomes, which, based on previous studies appears to be negative (i.e., very short sleep appears to be associated with poor health outcomes). Thus in effect, the row 3 regressions focus on the vastly predominant portion of our data. The fourth regression uses total sleep as an explanatory variable. The fifth regresses each health outcome on a dichotomous variable indicating the presence or absence of daytime sleep. Number of mobility limitations and the 1970s mortality risk score are significantly (positively) associated with age; right grip strength was significantly (negatively) associated with age (grip strength decreased by about 0.5 kg for each additional year of age); immediate recall was significantly associated with sex (females had lower recall than males) as was grip strength (lower by about 12.2 kg for females compared with males).

Table 4. Effects of objective and reported sleep durations on six health outcomes with adjustment for age and sex.

Moscow Pilot Project “Biological mechanisms of stress-related hazard in Russia”, 2002-03.

Explanatory variable, number of observations, and type of regression	Variable and type of regression					
	Mobility limitations, n=185	Right grip strength, n=181	Self-rated health, n=185	Cognitive recall, n=185	Stress Score, n=183	1970s mortality risk score. (Men only), n=144
	OLS	OLS	OLOGIT	OLS	OLS	OLS
1. Night sleep, reported	0.02 [0.05] [*] (0.777) [†]	-0.59 [0.37] (0.112)	0.04 [0.11] (0.699)	-0.18 [0.10] (0.081)	-0.01 [0.32] (0.979)	0.03 [0.03] (0.342)
2. Night sleep, objective	0.12 [0.06] (0.047)	-1.11 [0.37] (0.004)	0.18 [0.12] (0.127)	-0.26 [0.11] (0.018)	0.47 [0.33] (0.158)	0.05 [0.03] (0.115)
3. Night sleep, objective (if >5.5 h) [*]	0.15 [0.06] (0.032)	-1.51 [0.42] (0.000)	0.32 [0.13] (0.018)	-0.35 [0.12] (0.004)	0.65 [0.37] (0.087)	0.07 [0.03] (0.040)
4. Total sleep, objective	0.16 [0.06] (0.005)	-1.19 [0.35] (0.001)	0.22 [0.11] (0.040)	-0.25 [0.10] (0.015)	0.56 [0.07] (0.073)	0.03 [0.03] (0.231)
5. Presence of daytime sleep	0.59 [0.20] (0.003)	-2.54 [1.23] (0.041)	0.61 [0.38] (0.105)	-0.18 [0.36] (0.608)	1.77 [1.07] (0.102)	-0.07 [0.09] (0.446)

^{*} standard errors are given in square brackets. [†] p-values are given in parentheses.

[‡] Six observations with objective night sleep durations less than 5.5 hours are omitted.

CONCLUSION

It is recognized now that mortality is associated with longer sleep (4) even when controls are introduced for potentially confounding factors (3). Mechanisms underlying this important result remain largely unknown and the direction of the causal link is still disputable (44).

Our work provides the first piece of epidemiological evidence on the association between sleep and health outcomes among elderly Russians. It confirms the inverse relationship between duration of sleep and health in a high-mortality and high-stress context. The data suggest that in future studies, long sleep duration should be considered as a candidate risk factor of ill-health and death in Russia.

We also found also that sleep durations based on self-reports have the potential to significantly attenuate the relationships between sleep and health variables.

References

1. Kripke DF, Garfinkel L, Wingard DL, et al. Mortality associated with sleep duration and insomnia. *Archives of General Psychiatry* 2002;59:131-6.
2. Ayas NT, White DP, Manson JE, et al. A prospective study of sleep duration and coronary heart disease in women. *Archives of Internal Medicine* 2003;163:205-9.
3. Tamakoshi A, Ohno Y. Self-reported sleep duration as a predictor of all-cause mortality: results of JACC study, Japan. *Sleep* 2004; 27: 51-4.
4. Youngstedt, S, Kripke, D. Long sleep and mortality: rationale for sleep restriction. *Sleep Medicine* 2004;8:159-74.
5. Araujo AB, Yaggi HK, McKinlay JB. Sleep duration and risk of diabetes in middle-aged and older men: prospective results from the Massachusetts male aging study. *Annals of Epidemiology* 2004;14:598.
6. Wingard DL, Berkman LF. Mortality risk associated with sleep patterns among adults. *Sleep* 1983;6:102-7.
7. Qureshi AI, Giles WH, Croft JB, et al. Habitual sleep patterns and risk for stroke and coronary heart disease: a 10-year follow-up from NHANES I. *Neurology* 1997;48: 904-911.
8. Ohayon MM. Interactions between sleep normative data and sociocultural characteristics in the elderly. *Journal of Psychosomatic Research* 2004; 56:479-86.
9. Webb WB, Levy CM. Effects of spaced and repeated sleep deprivation. *Ergonomics* 1984;27:45-58.
10. Ferrara M, De Gennaro L, Casagrande M, et al. Selective slow-wave sleep deprivation and time-of-night effects on cognitive performance upon awakening. *Psychophysiology* 2000;37:440-6.
11. VanHelder T, Radomski MW. Sleep deprivation and the effect on exercise performance. *Sports Medicine (Auckland, N.Z.)* 1989;7:235-47.
12. Chen HI, Tang YR. Sleep loss impairs inspiratory muscle endurance. *The American Review of Respiratory Disease* 1989;140: 907-9.
13. Symons JD, Bell DG, Pope J, et al. Electro-mechanical response times and muscle strength after sleep deprivation. *Canadian Journal of Sport Sciences* 1988;13:225-30.
14. Trichopoulos D, Tzonou A, Christopoulos S, et al. Does a siesta protect from coronary heart disease? *The Lancet* 198;330:269-70.
15. Kalandidi A, Tzonou A, Toupadaki N, et al. A case-control study of coronary heart disease in Athens. *International Journal of Epidemiology* 1992;21:1074-80.
16. Campos H, Siles X. Siesta and the risk of coronary heart disease: results from a population-based, case-control study in Costa Rica. *International Journal of Epidemiology* 2000;29:429-37.
17. Bursztyn M, Ginsberg G, Hammerman-Rozenberg R, et al. The siesta in the elderly: risk factor for mortality? *Archives Of Internal Medicine* 1999;159:1582-6.
18. Burazeri G, Gofin J, Kark J. Over 8 hours of sleep – marker of increased mortality in Mediterranean population: follow-up population study. *Croatian Medical Journal* 2003;44:193-8.

19. Briones B, Adams N, Strauss M, et al. Relationship between sleepiness and general health status. *Sleep* 1996;19:583-8.
20. Newman AB, Spiekerman CF, Enright P, et al. Daytime sleepiness predicts mortality and cardiovascular disease in older adults. The Cardiovascular Health Study Research Group. *Journal of the American Geriatrics Society* 2000;48:115-23.
21. Foley D, Monjan A, Masaki K, et al. Daytime sleepiness is associated with 3-year incident dementia and cognitive decline in older Japanese-American men. *Journal of The American Geriatrics Society* 2001;49:1628-32.
22. McCall WV, Turpin E, Reboussin D, et al. Subjective estimates of sleep differ from polysomnographic measurements in obstructive sleep apnea patients. *Sleep* 1995;18:646-50.
23. Taheri S, Lin L, Austin D, et al. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLOS Medicine* 2004;1:e62. (www.plosmedicine.org)
24. Chervin RD, Guilleminault C. Overestimation of sleep latency by patients with suspected hypersomnolence. *Sleep* 1996;19:94-100.
25. Carscadon MA, Dement WC, Mitler MM, et al. Self-reports versus sleep laboratory findings in 122 drug-free subjects. *Am J Psychiatry* 1976;133:1382-88.
26. Gerardin J-L, Kripke D, Ancoli-Israel S, et al. Sleep duration, illumination, and activity patterns in a population sample: effects of gender and ethnicity. *Biol. Psychiatry* 2000;47:921-27.
27. Makarov L M, Shkol'nikova MA, Bereznitskaia VV, et al. Cardiac rhythm in sleeping healthy adolescents as recorded by Holter ECG monitoring. *Terapevticheskii Arkhiv* 1999;71:19-22.
28. Jurista F, van de Borne P, Migeotte P-F, et al. A study of the dynamic interactions between sleep EEG and heart rate variability in healthy young men. *Clinical Neurophysiology* 2003;114:2146-55.
29. Meslé F, Vallin J, Shkolnikov V. Reversal of mortality decline: the case of contemporary Russia. *World Health Statistics Quarterly* 2000;51:191-206.
30. Siegrist J. Place, social exchange and health: proposed sociological framework. *Social Science and Medicine* 2000;51:1283-93.
31. Heslop P, Davey Smith G, Metcalf C, et al. Sleep duration and mortality: the effect of short or long sleep duration on cardiovascular and all-cause mortality in working men and women. *Sleep Medicine* 2002; 3:305-14.
32. Deev AD, Shestov DB, Konstantinov VV. Population descriptions and methodology for US-USSR collaboration in Area 1 (Pathogenesis of Atherosclerosis): second prevalence study details for Moscow and Leningrad Lipid Research Clinics. *Atherosclerosis Reviews* 1988;17:103-109.
33. Dennis BH, Zhukovsky GS, Shestov DB, et al. The association of education with coronary heart disease mortality in the USSR Lipid Research Clinics Study. *International Journal of Epidemiology* 1993; 22:420-427.

34. Shestov DB, Deev AD, Klimov AN, et al. Increased risk of coronary heart disease death in men with low total and low-density lipoprotein cholesterol in the Russian lipid research clinics prevalence follow-up study. *Circulation* 1993; 88:846-53.
35. Deev A, Shkolnikov V. Neodnorodnost' smertnosti: analiz individualnikh dannyh. [Heterogeneity of mortality: analysis of individual-level data] In: Shkolnikov V, Andreev E, Maleva T. Eds. *Neravenstvo i Smertnost v Rossii [Inequality and Mortality in Russia]*. Carnegie Center, Moscow, 2000:63-83.
36. Plavinski SI, Plavinskaya SI, Klimov AN. Social factors and increase in mortality in Russia in the 1990s: prospective cohort study. *British Medical Journal* 2003;326:1240-42.
37. Shkolnikov VM, Deev AD, Kravdal Ø, et al. Educational differentials in male mortality in Russia and northern Europe. A comparison of an epidemiological cohort from Moscow and St. Petersburg with the male populations of Helsinki and Oslo. *Demographic Research* 2004;10:article 1. (www.demographic.research.org).
38. ACC/AHA. Guidelines for ambulatory electrocardiography: executive summary and recommendations. *Circulation* 1999;100:886-93.
39. McGue M, Christensen K. The heritability of level and rate-of-change in cognitive functioning in Danish twins aged 70 years and older. *Experimental Aging Research* 2002;28:435-51.
40. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *Journal of Health and Social Behavior* 1983;24:385-96.
41. Shkolnikova M, Shalnova S, Shkolnikov V, et al. Perceived stress, health, cognitive ability and their associations with heart rate variability and other physiological characteristics. Evidence from the Moscow Study on Biological Mechanisms of Stress-related Hazard. Presented at 25th Stress and Anxiety Research Society Conference, Book of Abstracts, University of Amsterdam, Amsterdam, 2004:215.
42. Marchioli R, Avanzini F, Barzi C et al. Assessment of absolute risk of death after myocardial infarction by use of multiple-risk-factor assessment equations. GISSI-Prevenzione mortality risk chart. *European Heart Journal* 2001;22:2085-103.
43. Stata Corporation. *Stata Reference Manual*. Release 7. College Station, Texas, 2001.
44. Foley DJ. An epidemiological perspective on one tale of a two-tailed hypothesis. *Sleep Medicine Reviews* 2004;8:155-7; discussion 175-176.
45. Nybo H, Gaist D, Jeune B, et al. Functional status and self-rated health in 2,262 nonagenarians: the Danish 1905 Cohort Survey. *Journal Of The American Geriatrics Society* 2001;49:601-9.
46. Idler ER, Benyamini Y. Self-rated health and mortality: a review of twenty-seven community studies. *Journal of Health and Social Behavior* 1997;38:21-37.
47. Opstad PK, Ekanger R, Nummerstad M, et. al. Performance, mood, and clinical symptoms in men exposed to prolonged, severe physical work and sleep deprivation. *Aviation, Space, and Environmental Medicine* 1978;49:1065-73.
48. Soldatos CR, Allaert FA, Ohta T, et al. How do individuals sleep around the world? Results from a single-day survey in ten countries. *Sleep Medicine* 2005;6:5-13.

49. Sajadieh A, Rasmussen V, Hein HO, et al. Familial predisposition to premature heart attack and reduced heart rate variability. *The American Journal of Cardiology* 2003; 92:234-6.
50. Pardo Y, Merz CN, Paul-Labrador M, et al. Heart rate variability reproducibility and stability using commercially available equipment in coronary artery disease with daily life myocardial ischemia. *The American Journal of Cardiology* 1996;78:866-70.
51. Guilleminault C, Connolly S, Winkle R, et al. A cyclical variation of the heart rate in sleep apnoea syndrome. Mechanisms, and usefulness of 24 h electrocardiography as a screening technique. *The Lancet* 1984;1:126-31.
52. Le Heuzey JY, Moutet JP, Guize L, et al. Measured and estimated sinoatrial conduction during variations in rhythm. Microelectrode study of the isolated rabbit atrium. *Archives Des Maladies Du Coeur Et Des Vaisseaux* 1984;77:146-54.
53. Roche F, Gaspoz JM, Court-Fortune I, et al. Screening of obstructive sleep apnea syndrome by heart rate variability analysis. *Circulation* 1999;100:1411-15.
54. Stein PK, Rich MW, Rottman JM, et al. Stability of index of heart rate variability in patients with congestive heart failure. *American Heart Journal* 1995;12:975-81.
55. Verrier RL, Muller JE, Hobson JA. Sleep, dreams, and sudden death: the case for sleep as an autonomic stress test for the heart. *Cardiovascular Research* 1996;31:181-211.
56. Lavery CE, Mittleman MA, Cohen MC, et al. Nonuniform nighttime distribution of acute cardiac events: a possible effect of sleep states. *Circulation* 1997; 96:3321-7.
57. Verrier RL. Sleep as an autonomic stress test for the heart. *Italian Heart Journal* 2000;1:329-30.
58. Versace F, Mozzato M, De Min Tona G, et al. Heart rate variability during sleep as a function of the sleep cycle. *Biological Psychology* 2003;63:149-62.